arXiv:1212.4221v1 [quant-ph] 18 Dec 2012

Photon induced tunneling in optomechanical systems

Xun-Wei Xu,[1] Yuan-Jie Li,[2] and Yu-xi Liu[1,3,*]

[1]*Institute of Microelectronics, Tsinghua University, Beijing 100084, China*
[2]*Department of Electronic Engineering, Tsinghua University, Beijing 100084, China*
[3]*Tsinghua National Laboratory for Information Science and Technology (TNList), Tsinghua University, Beijing 100084, China*

(Dated: December 19, 2012)

In contrast to recent studies [Rabl, Phys. Rev. Lett. 107, 063601 (2011); Nunnenkamp *et al.*, Phys. Rev. Lett. 107, 063602 (2011)] on photon blockade that prevents subsequent photons from resonantly entering the cavity in optomechanical systems, we study the photon induced tunneling that increases the probability of admitting subsequent photons in those systems. In particular, we analytically and numerically show how two- or three-photon tunneling can occur by avoiding single-photon blockade. Our study provides another way on photon control using a single mechanical resonator in optomechanical systems.

PACS numbers: 42.50.Wk, 07.10.Cm, 42.50.Ar, 42.65.-k

I. INTRODUCTION

Optomechanical systems [1, 2] have attracted extensive attention in the past few years because of its potential application in high-precision measurements and quantum information processing. To realize these benefits, the mechanical resonator has to be in its ground state; and the optomechanical radiation-pressure interaction strength should be bigger than the decay rates of the cavity field and the mechanical oscillator. The ground state cooling of the mechanical resonator has been experimentally studied in optomechanical systems (e.g., in Refs. [3–10]). Although the latter condition is not easy to be satisfied in standard optomechanical systems, however the experimentalists have obtained an effectively strong coupling, by applying a classical driving field to the cavity mode of the optomechanical systems, which has led to observations of normal-mode splitting (e.g., in Refs. [11, 12]) and optomechanically-induced transparency (e.g., in Refs. [13–15]). However, the resulting effective coupling resembles two linearly coupled harmonic oscillators [11] in their optical and mechanical position quadratures, and the coupling strength is proportional to the square root of the mean cavity photon number, thus it does not really describe the nonlinear effect at the single-photon level.

Recent two theoretical proposals [16, 17] showed that the single-photon effect or photon blockade [18] can occur when the optomechanical systems are approaching the strong coupling at the single-photon level [19–22]. This is because the frequency of the cavity field is parametrically modulated by the motion of the mechanical resonator in the optomechanical systems, this modulation results in the photon-photon interaction [23–28]. When the strong optomechanical radiation-pressure interaction makes the nonlinear photon-photon coupling strength bigger than the decay rate of the cavity field, then the photons can prevent the subsequent photons from resonantly entering the cavity. The photon blockade phenomenon has been demonstrated experimentally in cavity QED systems for microwave [29, 30] and optical [31, 32] photons. Meanwhile, the experiments [31, 32] also observed the photon induced tunneling, that is, the probability of admitting subsequent photons is increased when there is one photon inside the cavity. Moreover, absorption and emission of resonant photons in pairs have been observed in a single atom strongly coupled to an optical cavity [33, 34].

Motivated by these works [16–34] and also in contrast to the studies for photon blockade [16, 17], here we investigate photon-induced tunneling in the optomechanical systems. In Sec. II, we will describe the theoretical model of the system, and introduce the master equation, which is used to study statistical properties of the cavity field. The effect of the mechanical resonator on the mean photon numbers is discussed via the master equation when the optomechanical system reaches the steady state. In Sec. III, we will study the two-photon tunneling via the normalized second-order coherence function of the cavity field by numerically solving master equation. In Sec. IV, we further use the three-photon tunneling as an example to show how the multi-photon tunneling can happen. Finally, a summary is given in Sec. V.

II. THEORETICAL MODEL

We study an optomechanical system (see, review in Ref. [1]) in which a cavity field is coupled to a mechanical resonator through the radiation pressure. The Hamiltonian can be written as

$$H = \hbar\omega_0 a^\dagger a + \hbar\omega_m b^\dagger b + \hbar G a^\dagger a \left(b^\dagger + b\right), \qquad (1)$$

where a (b) and $a^\dagger$ ($b^\dagger$) are the annihilation and creation operators of the cavity field (mechanical resonator with the fundamental mode) with the frequency ω_0 (ω_m). The parameter G describes the coupling between the cavity field and the mechanical oscillator. To obtain the eigenvalues of the Hamiltonian in Eq. (1), let us apply a unitary transform $U = \exp\left[-Ga^\dagger a\left(b^\dagger - b\right)/\omega_m\right]$ to Eq. (1), and then we obtain an effective Hamiltonian $H' = UHU^\dagger$ with

$$H' = \hbar\omega_m b^\dagger b + \hbar\left(\omega_0 - \frac{G^2}{\omega_m}\right) a^\dagger a - \hbar\frac{G^2}{\omega_m} a^\dagger a^\dagger a a, \qquad (2)$$

---

[*] yuxiliu@tsinghua.edu.cn

which has eigenstates $|n, \widetilde{m}\rangle$ and corresponding eigenvalues

$$E_{n,m} = \hbar \left(n\,\omega_0 - n^2 \frac{G^2}{\omega_m} \right) + \hbar m\,\omega_m, \tag{3}$$

where $|n, \widetilde{m}\rangle \equiv U|n, m\rangle$, $|n, m\rangle$ represents that there are n photons and m phonons in the optomechanical system, respectively. Because a unitary transform does not change the eigenvalues of the Hamiltonian, thus the eigenvalues of the Hamiltonian in Eq. (1) are the same as in Eq. (3).

We now assume that the cavity field is coupled to a weak probe field with the frequency ω_c and the coupling strength ε_c. In the rotating reference frame with the unitary operator $R(t) = \exp[i\omega_c a^\dagger a t]$, the Hamiltonian in Eq. (1) becomes

$$\widetilde{H} = \hbar\Delta a^\dagger a + \hbar\omega_m b^\dagger b + \hbar G a^\dagger a \left(b^\dagger + b\right) + i\hbar\varepsilon_c \left(a^\dagger - a\right), \tag{4}$$

where $\Delta = \omega_0 - \omega_c$ is the detuning between the cavity field and the probe field. When the environmental effect is taken into account, the dynamical evolution of the reduced density operator ρ for the cavity field and the mechanical resonator can be described via the master equation [35]

$$\begin{aligned} \frac{\partial\rho}{\partial t} &= \frac{1}{i\hbar}\left[\widetilde{H}, \rho\right] + \frac{\gamma}{2}\left(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a\right) \\ &+ \frac{\gamma_m}{2}\left(2b\rho b^\dagger - b^\dagger b\rho - \rho b^\dagger b\right) \\ &+ \gamma_m \bar{n}_m \left(b\rho b^\dagger + b^\dagger \rho b - b^\dagger b\rho - \rho b b^\dagger\right), \end{aligned} \tag{5}$$

with the decay rates γ and γ_m of the cavity field and mechanical mode. $\bar{n}_m$ is the mean phonon number of the mechanical resonator given by the Bose-Einstein statistics $\bar{n}_m = [\exp(\hbar\omega_m/k_B T) - 1]^{-1}$ with the Boltzmann constant k_B and the environmental temperature T of the mechanical resonator. Here, the frequency of the cavity field is assumed to be high enough such that the thermal photon effect can be neglected. In the basis of the photon and phonon number states $|n, m\rangle$, the formal solution of the reduced density operator $\rho(t)$, governed by the master equation in Eq. (5), can be given by

$$\rho(t) = \sum_{n,m,n',m'} \rho_{n,m;n',m'}(t) |n, m\rangle\langle n', m'|. \tag{6}$$

If all elements $\rho_{n,m;n',m'}(t)$ in Eq. (6) are given, then any physical quantity of the optomechanical system can be calculated. For example, the mean photon number inside the cavity can be calculated as

$$\langle n\rangle = \left\langle a^\dagger a\right\rangle = \mathrm{Tr}[\rho(t) a^\dagger a] = \sum_{n,m} n\rho_{n,m;n,m}(t)\,. \tag{7}$$



FIG. 1: (Color online) The mean photon number $\langle n\rangle$ inside the cavity as a function of the normalized detuning Δ/Δ_0 in (a) for different decay rates of the mechanical mode at the given environmental temperature $T = 1\ \mu$K and (b) for different temperatures and given decay rate $\gamma_m/2\pi = 0.01$ MHz. Other parameters in (a) and (b) are $\varepsilon_c/2\pi = 0.01$ MHz, $G/2\pi = 2.5$ MHz, $\gamma/2\pi = 0.1$ MHz, and $\omega_m/2\pi = 10$ MHz.

Using Eqs. (5) and (7) and in the steady-state condition, the mean photon number $\langle n\rangle$ is plotted as a function of the normalized detuning Δ/Δ_0 with $\Delta_0 = G^2/\omega_m$ for: (i) different decay rates of the mechanical mode, e.g., $\gamma_m/2\pi = 0.01$ MHz, $\gamma_m/2\pi = 1$ MHz, and $\gamma_m/2\pi = 2$ MHz in Fig. 1 (a) and (ii) different environmental temperatures of the mechanical mode, e.g., $T = 1\ \mu$K, $T = 5$ mK, and $T = 10$ mK in Fig. 1 (b). We find that the mean photon number $\langle n\rangle$ is maximum at $(\Delta/\Delta_0) = 1$ which corresponds to the frequency $\omega_c = \omega_0 - (G^2/\omega_m)$ of the probe field. As shown in Eq. (2), this is because that the transition frequency of the cavity field from the one-photon state $|1\rangle$ to the vacuum (ground) state $|0\rangle$ is shifted to $\omega_0 - (G^2/\omega_m)$ when the mechanical resonator is coupled to the cavity field. Additionally, Fig. 1(a) and (b) show that when the decay rate γ_m or the environmental temperature T of the mechanical resonator increases, the full width at half maximum in the curve of the mean photon number $\langle n\rangle$ becomes broad for other given parameters. This means that the decay and environmental temeperature of the mechanical resonator significantly affect the lift time of the cavity photons.

III. TWO-PHOTON TUNNELING

The tunneling, blockade and other statistical properties of the photons can be characterized by the normalized nth-order correlation function at the zero time delay

$$g^{(n)}(0) = \frac{\left\langle a^{\dagger n} a^n\right\rangle}{\left\langle a^\dagger a\right\rangle^n} \equiv \frac{\mathrm{Tr}(\rho a^{\dagger n} a^n)}{\mathrm{Tr}(\rho a^\dagger a)^n}. \tag{8}$$

In our study here, the reduced density operator ρ of the cavity field and the mechanical resonator can be obtained by solving the master equation in Eq. (5).

Let us now study the simplest photon tunneling in pairs. Similar to the photon blockade, this phenomenon can be characterized by the normalized second-order correlation function $g^{(2)}(0)$ at the zero time delay. When $g^{(2)}(0)$ is less than one, the photons inside the cavity block subsequent photons from resonantly entering the cavity, and single-photons come out of the cavity. However, when $g^{(2)}(0)$ is bigger than one, photons inside the cavity enhance the resonantly entering probability of subsequent photons, this photon induced tunneling can make photons come out of the cavity in pairs under the certain condition. In Refs. [33, 36], the second-order differential correlation function at the zero time delay

$$C^{(2)}(0) = \left\langle a^{\dagger 2} a^2\right\rangle - \left\langle a^\dagger a\right\rangle^2 \equiv [g^{(2)}(0) - 1]\left\langle n\right\rangle^2, \tag{9}$$



FIG. 2: (Color online) $g^{(2)}(0)$ and $C^{(2)}(0)$ as functions of the normalized detuning Δ/Δ_0 and the coupling strength G for $T = 1\,\mu$K in (a) and (b), respectively. $g^{(2)}(0)$ as a function of Δ/Δ_0 for different temperatures T at $G/2\pi = 2.5$ MHz in (c). $C^{(2)}(0)$ as a function of Δ/Δ_0 for different temperatures T at $G/2\pi = 0.75$ MHz in (d). Other parameters are $\varepsilon_c/2\pi = 0.01$ MHz, $\gamma/2\pi = 0.1$ MHz, $\omega_m/2\pi = 10$ MHz, and $\gamma_m/2\pi = 0.01$ MHz.

is introduced to characterize the probability of creating photon pairs simultaneously in the cavity, where $\langle n\rangle$ is the mean photon number inside the cavity.

To explore two-photon tunneling and compare it with the photon blockade, we study $g^{(2)}(0)$ and $C^{(2)}(0)$ in the steady-state. In Figs. 2(a) and (b), $g^{(2)}(0)$ and $C^{(2)}(0)$ versus the detuning $\Delta = \omega_0 - \omega_c$ and the coupling strength G are plotted at the environmental temperatures $T = 1\,\mu$K of the mechanical resonator. In each figure, there are two curves which correspond to single-photon (green dashed curve for $\Delta = \Delta_0$) and two-photon (black dashed curve $\Delta = 2\Delta_0$) resonant excitations from the ground state to the first excited and the second excited states of the cavity field, respectively. Figs. 2(a) and (b) clearly shows $g^{(2)}(0) < 1$ and $C^{(2)}(0) < 0$ for the single-photon resonant excitation from the ground state to the first excited state when the coupling strength G is bigger than the decay rate γ of the cavity field, that is, the single-photon phenomena or photon blockade occurs as shown in Refs. [16, 17]. However, if the frequency of the probe field equals to the half of the transition frequency from the ground state to the second excited state, i.e., $\Delta = 2\Delta_0$, then the single-photon transition from the ground state to the first excited state is suppressed, but the second photon can enter the cavity and makes resonant transition from the ground state to the second excited state together with the first photon. This photon-induced tunneling means that the cavity can absorb photons in pairs simultaneously, therefore we can find $g^{(2)}(0) > 1$ and $C^{(2)}(0) > 0$ in Figs. 2(a) and (b).

We further showed the effect of environmental temperature on the photon tunneling via $g^{(2)}(0)$ and $C^{(2)}(0)$ in Figs. 2(c) and (d). We find that the low temperature can make photon-induced tunneling and photon blockade easy to be observed, because with the decrease of the temperature, $g^{(2)}(0)$ and $C^{(2)}(0)$ decrease under the single-photon ($\Delta = \Delta_0$) resonant driving, and increase under the two-photon ($\Delta = 2\Delta_0$) resonant driving. As shown in Fig. 3(a), there is an additional resonant peak for $g^{(2)}(0)$ at $\Delta = 3\Delta_0$, which corresponds to the transition between two eigenstates $|1, \widetilde{m}\rangle$ and $|2, \widetilde{m}\rangle$. We find that the height of the resonant peak at $\Delta = 3\Delta_0$ increases when the temperature becomes higher, qualitatively, this is because the environmental temperature can change the population distribution, especially it can enhance population distribution in higher energy levels, and then the transition from $|1, \widetilde{m}\rangle$ to $|2, \widetilde{m}\rangle$ is also enhanced. All these result in the increase of $g^{(2)}(0)$ at $\Delta = 3\Delta_0$ when the temperature becomes higher.



FIG. 3: (Color online) (a) Energy level diagram for the optomechanical system. Here, the horizontal long black lines linked by the vertical black lines denote the energy levels of the cavity field with frequency ω_0. The parameter ω_c is the frequency of the driven field. $\Delta = \omega_0 - \omega_c$. The horizontal bright purple lines denote the energy levels ($|n, \widetilde{m}\rangle$) of the optomechanical system with $\Delta_0 = G^2/\omega_m$. The horizontal long blue lines linked by the vertical blue line with two arrows denotes the transition of $|1, \widetilde{m}\rangle \rightarrow |2, \widetilde{m}\rangle$. (b) $g^{(2)}(0)$ is plotted as functions of G/ω_m for $\Delta = \Delta_0$, $T = 1\,\mu$K. (c) $g^{(2)}(0)$ is plotted as functions of G/ω_m for $\Delta = 0$, $T = 1\,\mu$K (black) and 2 mK (red). The other parameters are the same as in Fig. 2(c).

The effect of the phonon states on the photon blockade has already been studied in Refs. [16, 27]. It was found that under the photon blockade condition $\Delta = \Delta_0$, the two-photon resonant transition between $|0, \widetilde{0}\rangle$ and $|2, \widetilde{m}\rangle$ can also occur if $\Delta_0 = m\omega_m/2$. Thus there are resonant peaks located at $G/\omega_m = \sqrt{m/2}$ for $g^{(2)}(0)$ as shown in Fig. 3(b) for different phonon states $|m\rangle$ (e.g., $m = 1, 2, 3, 4$). The phonon states can also affect the resonant condition of the photon-induced tunneling. As shown in Fig. 3(c), under the resonant condition $\Delta = 0$, there are five resonant peaks for $g^{(2)}(0) \gg 1$ with $G < \omega_m$. As shown Fig. 3(a), these five peaks correspond to two types of the resonant conditions: (i) if Δ_0 satisfies the condition $4\Delta_0 = m\omega_m$, then two-photon resonant excitation between $|0, \widetilde{0}\rangle$ and $|2, \widetilde{m}\rangle$ can occur with the peaks located at $G/\omega_m = \sqrt{m/4}, m = 1, 2, 3$; (ii) if Δ_0

satisfies the condition $3\Delta_0 = m\omega_m$, then resonant excitation from $|1, \widetilde{0}\rangle$ to $|2, \widetilde{m}\rangle$ is allowed with the peaks located at $G/\omega_m = \sqrt{m/3}, m = 1, 2$. We also find that the transition of $|1, \widetilde{0}\rangle \rightarrow |2, \widetilde{m}\rangle$ is enhanced with the increase of the temperature, so the resonant peak at $G/\omega_m = \sqrt{m/3}, m = 1, 2$ increases as the temperature becomes higher.

To further analyze the conditions of the photon blockade and the photon induced tunneling, we now study the relation between $g^{(2)}(0)$ and the probabilities $P(n)$ of n-photon distribution. Because the probabilities $P(n)$ corresponding to the state in Eq. (6) are $P(n) = \sum_m \rho_{n,m;n,m}(t)$, then we have

$$g^{(2)}(0) = \frac{\mathrm{T}r[\rho a^{\dagger 2} a^2]}{[\mathrm{T}r(\rho a^\dagger a)]^2} = \frac{\sum\limits_n n(n-1)P(n)}{\left[\sum\limits_n nP(n)\right]^2}. \tag{10}$$

In the limit of the weak probe field, e.g., $\varepsilon_c = 0.1\gamma$, $P(n) \gg P(n+1)$ and in the steady-state, $g^{(2)}(0)$ can be given approximately as

$$g^{(2)}(0) \approx \frac{2P(2)}{[P(1)]^2} \approx 4\left|\frac{\frac{\gamma}{2} + i(\Delta - \Delta_0)}{\gamma + i2(\Delta - 2\Delta_0)}\right|^2, \tag{11}$$

here we have assumed that the phonon is in its ground state to obtain second approximated expression. This assumption has also been made in following derivations in Eq. (15) and Eq. (17). Eq. (11) means that $g^{(2)}(0)$ is proportional to the ratio between the probability to prepare a two-photon state and that to prepare two single-photon states independently. In Fig. 4(a), $g^{(2)}(0)$ and its approximated expression in Eq. (11) versus the normalized detuning Δ are numerically simulated in the steady-state. We find that the approximated expression fits well with the exact solution of $g^{(2)}(0)$ in the limit of the weak probe field. Therefore, $g^{(2)}(0) > 1 (g^{(2)}(0) < 1)$ means that the probability to excite two-photon state is bigger (smaller) than that to excite two single-photon states independently, and then photon-induced tunneling (photon blockade) happens. To further explore the relation between the height of Fig. 2(c) (and also Fig. 4(a)) and system parameters, from Eq. (11), we can approximately obtain

$$g^{(2)}(0) \approx \frac{\gamma^2 + 4\Delta_0^2}{\gamma^2}, \tag{12}$$

for $\Delta = 2\Delta_0$, which corresponds to the height of the peak in Fig. 4 (a) and Fig. 2(c) at $\Delta = 2\Delta_0$. Thus the height of the peak in Fig. 4(a) is approximately determined by the ratio between the decay rate γ of the cavity field and the parameter Δ_0. We can approximately obtain another formula

$$g^{(2)}(0) \approx \frac{\gamma^2}{\gamma^2 + 4\Delta_0^2}, \tag{13}$$

for $\Delta = \Delta_0$, which corresponds to the deep in Fig. 4 (a) and Fig. 2(c) at $\Delta = \Delta_0$. It is clear that the conditions to observe photon blockade or photon induced tunneling respectively are

$$4\Delta_0^2 \gg \gamma^2 \quad \text{or} \quad \Delta_0 \gg \gamma/2. \tag{14}$$



FIG. 4: (Color online) (a) $g^{(2)}(0)$ and $2P(2)/P(1)^2$, (b) $g^{(3)}(0)$, $6P(3)/P(1)^3$, $g_2^{(3)}(0)$ and $3P(3)/[P(1)P(2)]$ plotted as functions of the normalized detuning Δ/Δ_0 for $G = 2.5$ MHz and $T = 1\,\mu$K. The other parameters are the same as in Fig. 2(c).

IV. MULTI-PHOTON TUNNELING

In principle, the multi-photon tunneling (more than two) can be studied via Eq. (8). However, for the sake of simplicity, we just use the normalized third-order correlation function $g^{(3)}(0)$ as an example to show multi-photon tunneling. Using the same discussions as for Eq. (10) and Eq. (11), $g^{(3)}(0)$ can be approximately written as

$$\begin{aligned} g^{(3)}(0) &\equiv \frac{\mathrm{T}r\left(\rho a^{\dagger 3} a^3\right)}{[\mathrm{T}r(\rho a^\dagger a)]^3} \approx \frac{6P(3)}{[P(1)]^3} \\ &\approx \left|\frac{6\left[\frac{\gamma}{2} + i(\Delta - \Delta_0)\right]^2}{[\gamma + i2(\Delta - 2\Delta_0)]\left[\frac{3}{2}\gamma + i3(\Delta - 3\Delta_0)\right]}\right|^2 \end{aligned} \tag{15}$$

in the steady-state case with the weak probe field. It is clear that $g^{(3)}(0)$ is proportional to the ratio between the probability to create a three-photon state and that to create three single-photons independently. If we introduce a quantity

$$g_2^{(3)}(0) \equiv \frac{g^{(3)}(0)}{g^{(2)}(0)} = \frac{\mathrm{T}r\left(\rho a^{\dagger 3} a^3\right)}{\mathrm{T}r(\rho a^\dagger a)\,\mathrm{T}r\left(\rho a^{\dagger 2} a^2\right)} \tag{16}$$

to characterize the ratio between the normalized third-order and second-order correlation functions, then in the steady-state case, we can approximately obtain

$$g_2^{(3)}(0) \approx \frac{3P(3)}{P(1)P(2)} \approx 9\left|\frac{\frac{\gamma}{2} + i(\Delta - \Delta_0)}{\frac{3}{2}\gamma + i3(\Delta - 3\Delta_0)}\right|^2 \tag{17}$$

which is proportional to the ratio between the probability to create a three-photon state and the joint probability to create a single-photon and a two-photon state independently, which is also a possible way to create three photons.

In Fig. 4(b), $g^{(3)}(0)$ and its approximated expression $6P(3)/[P(1)]^3$ as well as $g_2^{(3)}(0)$ and its approximated expression $3P(3)/[P(1)P(2)]$ are plotted as functions of Δ/Δ_0. We find that the approximation in both Eq. (15) and Eq. (17) is valid for the weak probe field. Under the condition that $4\Delta_0^2 \gg \gamma^2$, we can further find:

$$g^{(3)}(0) \approx \left(\frac{\gamma^2}{8\Delta_0^2}\right)^2, \quad g_2^{(3)}(0) \approx \frac{\gamma^2}{16\Delta_0^2}, \tag{18}$$

for the resonant condition $\Delta = \Delta_0$; and

$$g^{(3)}(0) \approx \frac{4\Delta_0^2}{\gamma^2}, \quad g_2^{(3)}(0) \approx 1, \tag{19}$$

for the resonant condition $\Delta = 2\Delta_0$; as well as

$$g^{(3)}(0) \approx \frac{64\Delta_0^2}{\gamma^2}, \quad g_2^{(3)}(0) \approx \frac{16\Delta_0^2}{\gamma^2}, \tag{20}$$

for the resonant condition $\Delta = 3\Delta_0$. Eqs. (19-20) show that the heights of the resonant peaks are approximately determined by the ratio of Δ_0 and γ as for two-photon tunneling.

Fig. 4(b) and Eqs. (18-20) show that: (i) for the single-photon resonant excitation ($\Delta = \Delta_0$), $g^{(3)}(0) < 1$ and $g_2^{(3)}(0) < 1$, this tells us that the photon inside the cavity is antibunching or the subsequent photons will be blocked by the photons inside the cavity. (ii) For the three-photon resonant excitation ($\Delta = 3\Delta_0$), $g^{(3)}(0) > 1$ and $g_2^{(3)}(0) > 1$, that is, the photon inside the cavity is bunching or the cavity can absorb three photons simultaneously. (iii) Remarkably, for the two-photon resonant excitation ($\Delta = 2\Delta_0$), $g^{(3)}(0) > 1$ which means that the probability of three-photon absorption is bigger than that of three single-photon absorption independently. We also find $g_2^{(3)}(0) \approx 1$ at the point $\Delta = 2\Delta_0$. This tells us an expected result that the joint probability of the photon absorption by the cavity in pairs after or before single-photon absorption is approximately equal to that of three-photon absorption by the cavity. Therefore, the necessary condition for absorbing three photons by the cavity simultaneously (or three-photon tunneling) is $\Delta = 3\Delta_0$. In this condition, $g^{(3)}(0) > 1$ and $g_2^{(3)}(0) > 1$, the probability of generating three photons simultaneously is bigger than those of generating three single-photons independently and generating a single-photon after or before generating a photon pair.

V. CONCLUSIONS

In summary, we have studied the photon induced tunneling and compare it with the photon blockade in optomechanical systems with the single-photon strong coupling condition. Our study shows that the cavity field can exhibit photon antibunching when the probe field is resonant with the transition from the ground state to the first excited state of the cavity field. However, the two-photon tunneling occurs when the frequency of the driving field equals to the half of the transition frequency from the ground state to the second excited state of the cavity field. Moreover, we also find that three-photon tunneling occurs when the frequency of the probe field satisfies the condition of the three-photon resonant excitation from the ground state to the third excited state. Our studies also show that the phonon states greatly affect the multi-photon resonance in the strong optomechanical coupling at the single-photon level. These studies can be easily generalized to the multi-photon tunneling case. Our results show that it is possible to observe photon induced tunneling in the optomechanical systems under the single-photon strong coupling condition.

VI. ACKNOWLEDGEMENT

YXL is supported by the National Natural Science Foundation of China under Nos. 10975080 and 61025022.

---

[1] T. J. Kippenberg and K. J. Vahala, Science **321**, 1172 (2008), and the references therein.
[2] F. Marquardt and S. M. Girvin, Physics **2**, 40 (2009).
[3] O. Arcizet, P.-F. Cohadon, T. Briant, M. Pinard, and A. Heidmann, Nature **444**, 71 (2006).
[4] D. Kleckner and D. Bouwmeester, Nature **444**, 75 (2006).
[5] S. Gigan, H. R. Böhm, M. Paternostro, F. Blaser, G. Langer, J. B. Hertzberg, K. C. Schwab, D. Bäuerle, M. Aspelmeyer, and A. Zeilinger, Nature **444**, 67 (2006).
[6] A. Schliesser, P. Del'Haye, N. Nooshi, K. J. Vahala, and T. J. Kippenberg, Phys. Rev. Lett. **97**, 243905 (2006).
[7] J. D. Teufel, J. W. Harlow, C. A. Regal, and K. W. Lehnert, Phys. Rev. Lett. **101**, 197203 (2008).
[8] J. D. Thompson, B. M. Zwickl, A. M. Jayich, Florian Marquardt, S. M. Girvin, and J. G. E. Harris, Nature **452**, 72 (2008).
[9] T. Rocheleau, T. Ndukum, C. Macklin, J. Hertzberg, A. Clerk, and K. Schwab, Nature **463**, 72 (2009).
[10] J. D. Teufel, T. Donner, Dale Li, J. W. Harlow, M. S. Allman, K. Cicak, A. J. Sirois, J. D. Whittaker, K. W. Lehnert, and R. W. Simmonds, Nature **475**, 359 (2009).
[11] S. Groblacher, K. Hammerer, M. R. Vanner, and M. Aspelmeyer, Nature **460**, 724 (2009).
[12] J. D. Teufel, Dale Li, M. S. Allman, K. Cicak, A. J. Sirois, J. D. Whittaker, and R. W. Simmonds, Nature **471**, 204 (2011).
[13] G. S. Agarwal and Sumei Huang, Phys. Rev. A **81**, 041803(R) (2010).
[14] S. Weis, R. Riviere, S. Deleglise, E. Gavartin, O. Arcizet, A. Schliesser, and T. J. Kippenberg, Science **330**, 1520 (2010).
[15] A. H. Safavi-Naeini, T. P. Mayer Alegre, J. Chan, M. Eichenfield, M. Winger, Q. Lin, J. T. Hill, D. E. Chang, and O. Painter, Nature **472**, 69 (2011).
[16] P. Rabl, Phys. Rev. Lett. **107**, 063601 (2011).
[17] A. Nunnenkamp, K. Børkje, and S. M. Girvin, Phys. Rev. Lett. **107**, 063602 (2011).
[18] A. Imamoğlu, H. Schmidt, G. Woods, and M. Deutsch, Phys. Rev. Lett. **79**, 1467 (1997).
[19] S. Gupta, K. L. Moore, K. W. Murch, and D. M. Stamper-Kurn, Phys. Rev. Lett. 99, 213601 (2007).
[20] F. Brennecke, S. Ritter, T. Donner and T. Esslinger, Science **322**, 235 (2008).
[21] M. Eichenfield, J. Chan, R. M. Camacho, K. J. Vahala, and O. Painter, Nature **462**, 78 (2009).
[22] J. D. Teufel, D. Li, M. S. Allman, K. Cicak, A. J. Sirois, J. D. Whittaker, and R. W. Simmonds, Nature **471**, 204 (2011).
[23] P. Meystre, E. M. Wright, J. D. McCullen, and E. Vignes, J. Opt. Soc. Am. B **2**, 1830 (1985).
[24] S. Mancini and P. Tombesi, Phys. Rev. A **49**, 4055 (1994).
[25] Z. R. Gong, H. Ian, Y. X. Liu, C. P. Sun, and F. Nori, Phys. Rev.

A **80**, 065801 (2009).

[26] J. Q. Liao, H. K. Cheung, and C. K. Law, Phys. Rev. A **85**, 025803 (2012).

[27] J. Q. Liao and C. K. Law, arXiv:1206.3085.

[28] B. He, Phys. Rev. A **85**, 063820 (2012).

[29] C. Lang, D. Bozyigit, C. Eichler, L. Steffen, J. M. Fink, A. A. Abdumalikov, Jr., M. Baur, S. Filipp, M. P. da Silva, A. Blais, and A. Wallraff, Phys. Rev. Lett. **106**, 243601 (2011).

[30] A. J. Hoffman, S. J. Srinivasan, S. Schmidt, L. Spietz, J. Aumentado, H. E. Türeci, and A. A. Houck, Phys. Rev. Lett. **107**, 053602 (2011).

[31] K. M. Birnbaum, A. Boca, R. Miller, A. D. Boozer, T. E. Northup, and H. J. Kimble, Nature **436**, 87 (2005).

[32] A. Faraon, I. Fushman, D. Englund, N. Stoltz, P. Petro, and J. Vuckovic, Nature Phys. **4**, 859 (2008).

[33] A. Kubanek, A. Ourjoumtsev, I. Schuster, M. Koch, P. W. H. Pinkse, K. Murr, and G. Rempe, Phys. Rev. Lett. **101**, 203602 (2008).

[34] M. Koch, C. Sames, M. Balbach, H. Chibani, A. Kubanek, K. Murr, T. Wilk, and G. Rempe, Phys. Rev. Lett. **107**, 023601 (2011).

[35] H. J. Carmichael, *An Open Systems Approach to Quantum Optics*, (Springer-Verlag, Berlin, 1993).

[36] A. Majumdar, M. Bajcsy, and J. Vuckovic, arXiv:1106.1926.